


# Use Of Medical Imaging Measures - State

State	Outpatients with low back pain who had an MRI without trying recommended treatments first, such as physical therapy.	
AK		38.7
AL		34.9
AR		37.4
AS	N/A	
AZ		30.9
CA		28.4
CO		35.8
CT		26.9
DC		31.9
DE		25.8
FL		28.3
GA		31
GU	N/A	
HI		30.1
IA		30.2
ID		37.3
IL		33.4
IN		35.2
KS		34.5

# Use Of Medical Imaging Measures - State




# Use Of Medical Imaging Measures - State

Outpatient CT scans of the abdomen that were “combination” (double) scans.

	0.081
	0.258
	0.208
N/A	
	0.167
	0.228
	0.163
	0.12
	0.126
	0.258
	0.219
	0.151
N/A	
	0.114
	0.163
	0.121
	0.246
	0.237
	0.365

# Use Of Medical Imaging Measures - State

Outpatient CT scans of the chest that were “combination” (double) scans.



# Use Of Medical Imaging Measures - State

KY		35.6
LA		36.8
MA		30.1
MD		31.7
ME		36.7
MI		32
MN		39.7
MO		36
MP	N/A	
MS		37.1
MT		37.3
NC		30.6
ND		31.9
NE		33.8
NH		31.5
NJ		26.4
NM		33.9
NV		33.2
NY		26.9
OH		30.2
OK		34.3
OR		36.2

# Use Of Medical Imaging Measures - State



# Use Of Medical Imaging Measures - State

	0.16
	0.36
	0.109
	0.064
	0.133
	0.135
	0.1
	0.227
N/A	
	0.203
	0.181
	0.115
	0.177
	0.152
	0.078
	0.126
	0.142
	0.178
	0.141
	0.164
	0.372
	0.115


# Use Of Medical Imaging Measures - State

	0.054
	0.087
	0.01
	0.013
	0.045
	0.068
	0.018
	0.059
N/A	
	0.075
	0.039
	0.037
	0.025
	0.033
	0.014
	0.026
	0.057
	0.047
	0.024
	0.038
	0.134
	0.022


# Use Of Medical Imaging Measures - State

PA		32.4
PR		31.7
RI		31.8
SC		34.4
SD		31.3
TN		30.7
TX		34.7
UT		34.8
VA		31.5
VI	N/A	
VT		27.1
WA		35.6
WI		31.8
WV		38.5
WY		40.8

# Use Of Medical Imaging Measures - State



# Use Of Medical Imaging Measures - State



# Use Of Medical Imaging Measures - State

